

FOXD4 blocking peptide

Blocking peptide binds specifically to the target antibody and block undesirable antibody binding. This occurs because the peptide resembles the epitope for which the antibody is specific, therefore, the antibody binds to the blocking peptide are no longer able to bind to the epitope in the protein.

Pre-adsorption with blocking peptide greatly improves the results and it is extremely useful in immunohistochemistry (IHC) or in Western blotting (immunobloting) when non-specific binding is an issue.

This peptide is recommended to block anti-FOXD4 Antibody (Catalog no. 4075498) reactivity

Directions for Use: Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours

| Catalog No. | 5075498 |
|------------------|--|
| Size | lmg |
| Product Category | Blocking Peptide |
| Gene Symbol | FOXD4 |
| UniProt ID | Q12950; Q8WXT5; Q5VV16; Q6VB85; Q6VB84; Q3SYB3 |
| Purity | >85% |

| Synonyms | FOXD4; FKHL9; FOXD4A; FREAC5; Forkhead box protein D4; Forkhead- |
|-------------------|--|
| | related protein FKHL9; Forkhead-related transcription factor 5; FREAC-5; |
| | Myeloid factor-alpha; FOXD4L4; FOXD4B; Forkhead box protein D4-like 4; |
| | FOXD4-like 4; Forkhead box protein D4B; Myeloid factor-gamma; |
| | FOXD4L5; Forkhead box protein D4-like 5; FOXD4-like 5; FOXD4L2; |
| | Forkhead box protein D4-like 2; FOXD4-like 2; FOXD4L3; Forkhead box |
| | protein D4-like 3; FOXD4-like 3; FOXD4L6; Forkhead box protein D4-like |
| | 6; FOXD4-like 6 |
| Storage/Stability | -20°C/1 year |
| Shipping | Gel Packs |

www.realgenelabs.com

For Research Use only