

MED28 blocking peptide

Blocking peptide binds specifically to the target antibody and block undesirable antibody binding. This occurs because the peptide resembles the epitope for which the antibody is specific, therefore, the antibody binds to the blocking peptide are no longer able to bind to the epitope in the protein.

Pre-adsorption with blocking peptide greatly improves the results and it is extremely useful in immunohistochemistry (IHC) or in Western blotting (immunoblotting) when non-specific binding is an issue.

This peptide is recommended to block anti-MED28 Antibody (Catalog no. 4073656) reactivity

Directions for Use: Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours

Catalog No.	5073656
Size	1mg
Product Category	Blocking Peptide
Gene Symbol	MED28
UniProt ID	Q9H204
Purity	>85%
Synonyms	EG1; Mediator of RNA polymerase II transcription subunit 28; Endothelial-derived protein 1; Mediator complex subunit 28; Merlin and Grb2-interacting cytoskeletal protein; Magicin; Tumor angiogenesis marker EG-1
Storage/Stability	-20°C/1 year

Shipping

Gel Packs

www.realgenelabs.com

For Research Use only