

MUC1 (pY1229) blocking peptide

Blocking peptide binds specifically to the target antibody and block undesirable antibody binding. This occurs because the peptide resembles the epitope for which the antibody is specific, therefore, the antibody binds to the blocking peptide are no longer able to bind to the epitope in the protein.

Pre-adsorption with blocking peptide greatly improves the results and it is extremely useful in immunohistochemistry (IHC) or in Western blotting (immunobloting) when non-specific binding is an issue.

This peptide is recommended to block anti-MUC1 (pY1229) Antibody (Catalog no. 4071772) reactivity

Directions for Use: Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours

Catalog No.	5171772
Size	1mg
Product Category	Blocking Peptide
Gene Symbol	MUC1
UniProt ID	P15941
Purity	>85%
Synonyms	PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Cancer antigen 15-3; CA 15-3; Carcinoma-associated mucin; Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic epithelial mucin; PEM; Tumor-associated epithelial membrane antigen; EMA; Tumor-associated mucin; CD227

Storage/Stability	-20°C/1 year	
Shipping	Gel Packs	

www.realgenelabs.com

For Research Use only