

## Vasopressin V2 Receptor blocking peptide

---

Blocking peptide binds specifically to the target antibody and block undesirable antibody binding. This occurs because the peptide resembles the epitope for which the antibody is specific, therefore, the antibody binds to the blocking peptide are no longer able to bind to the epitope in the protein.

Pre-adsorption with blocking peptide greatly improves the results and it is extremely useful in immunohistochemistry (IHC) or in Western blotting (immunoblotting) when non-specific binding is an issue.

This peptide is recommended to block anti-Vasopressin V2 Receptor Antibody (Catalog no. 4074578) reactivity

Directions for Use: Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours

Catalog No.	5074578
Size	1mg
Product Category	Blocking Peptide
Gene Symbol	AVPR2
UniProt ID	P30518
Purity	>85%
Synonyms	ADHR; DIR; DIR3; V2R; Vasopressin V2 receptor; V2R; AVPR V2; Antidiuretic hormone receptor; Renal-type arginine vasopressin receptor
Storage/Stability	-20°C/1 year

Shipping

Gel Packs

[www.realgenelabs.com](http://www.realgenelabs.com)

For Research Use only